

Measures and semi-orthogonal functions on the unit circle

M.J.Cantero, M.P.Ferrer, L.Moral, L.Velázquez

Departamento de Matemática Aplicada. Universidad de Zaragoza. Spain.

Abstract.

The zeros of semi-orthogonal functions with respect to a probability measure μ supported on the unit circle can be applied to obtain Szegő quadrature formulas. The discrete measures generated by these formulas weakly converge to the orthogonality measure μ . In this paper we construct families of semi-orthogonal functions with interlacing zeros, and give a representation of the support of μ in terms of the asymptotic distribution of such zeros.

Keywords and phrases: Quadrature formulas, Semi-orthogonal functions, Asymptotic distribution of zeros, Support of a measure.

(1991) *AMS Mathematics Subject Classification* : 42C05

§ 1 - Introduction.

Given a probability measure ν on the real line, it is well known (see for example [5] or [4]) that a sequence of orthogonal polynomials (SOP), $(p_n)_{n \in \mathbb{N}}$, with respect to ν , satisfies the following properties:

- a) The zeros of p_n are real and simple for $n \geq 1$.
- b) If I is an interval such that $\text{supp } \nu \subseteq I$, the zeros of p_n lie on the interior of I for $n \geq 1$.
- c) The polynomials p_n and p_{n+1} have interlacing zeros for $n \geq 1$.
- d) The interpolatory quadrature formulas

$$\int_{\mathbb{R}} F(x) d\nu(x) \approx \sum_{k=1}^n F(x_k^{(n)}) H_k^{(n)}, \quad n \geq 1,$$

are exact for every polynomial F with degree not greater than $2n - 1$ if the nodes $(x_k^{(n)})_{k=1}^n$ are the zeros of p_n and $H_k^{(n)} = 1/K_{n-1}(x_k^{(n)}, x_k^{(n)})$, where

$K_n(x, y)$ is the n -kernel associated to $(p_n)_{n \in \mathbb{N}}$, which is

$$K_n(x, y) = \frac{k_n}{k_{n+1}} \frac{p_{n+1}(x)p_n(y) - p_n(x)p_{n+1}(y)}{(x - y) \int_{\mathbb{R}} p_n^2(t) d\nu(t)},$$

being k_n the leading coefficient of p_n . These quadrature formulas imply the existence of a sequence $(\nu_n)_{n \geq 1}$ of discrete measures on the real line given by

$$d\nu_n(x) = \sum_{k=1}^n \frac{\delta(x - x_k^{(n)})}{K_{n-1}(x_k^{(n)}, x_k^{(n)})} dx$$

(δ is the Dirac distribution), such that $\nu_n \xrightarrow{*} \nu$ when the support of ν is bounded.

Therefore, there is a close relationship between the measure ν and the location of the zeros of a SOP. In fact, properties a), b), c), d) can be used to study the support of ν in terms of the asymptotics of such zeros (see [5], [4]).

As for measures on the unit circle $\mathbb{T}$, the situation is rather different. The associated orthogonal polynomials have their zeros inside the open unit disk, and not on $\mathbb{T}$, where the measure is supported. Moreover, these zeros are not necessarily simple. So, it has no meaning to look for results completely analogous to the ones above.

In the case of the real line, we can find other functions that satisfy property a), and a weak version of b), c), d): polynomials $q_n(x)$, $n \geq 1$, with $\deg(q_n) = n$, and orthogonal to x^k for all $k < n - 1$, but not to x^{n-1} . These are called quasi-orthogonal polynomials (see [5], [4]), and the modified versions of properties b), c), d) that they satisfy are:

- b') If I is an interval such that $\text{supp } \nu \subseteq I$, it is not possible to ensure that all the zeros of q_n lie on the interior of I , but at most one of them lies outside.
- c') It is not always true that q_n and q_{n+1} have interlacing zeros, but, if the convex hull of $\text{supp } \nu$ is not $\mathbb{R}$, the sequence $(q_n)_{n \geq 1}$ can be chosen so that all the polynomials q_n have a common zero outside this convex hull, and the remaining zeros of q_n and q_{n+1} (that are inside this convex hull) interlace.
- d') The interpolatory quadrature formulas constructed by using as nodes the zeros of q_n are exact for every polynomial with degree not greater than $2n - 2$, instead of $2n - 1$. That is, they are exact on a $(2n - 1)$ -dimensional vector subspace of the space of real polynomials. Likewise, these quadrature formulas generate a sequence of discrete measures on the real line weakly converging to ν .

As in the case of orthogonal polynomials, properties a), b'), c'), d') allow to study the support of ν through the analysis of the asymptotic distribution of zeros of quasi-orthogonal polynomials.

The aim of this paper is to get on $\mathbb{T}$ a situation as similar as possible to the one we have in the real line with quasi-orthogonal polynomials: given a probability measure μ on $\mathbb{T}$, we look for sequences $(F_n)_{n \geq 1}$ of analytic functions on a open subset $\Omega \supset \mathbb{T}$ of $\mathbb{C}$, such that, when restricted to $\mathbb{T}$ can be viewed as real functions, and that satisfy analogous properties to a), b'), c'), d'). That is,

1. $F_n(z)$ is real for $z \in \mathbb{T}$.
2. The zeros of F_n are simple and lie on $\mathbb{T}$.
3. If I is a connected subset of $\mathbb{T}$ such that $\text{supp } \mu \subseteq I$, at most one zero of F_n lies on the closure of $\mathbb{T} \setminus I$.
4. The zeros of the functions F_n and F_{n+1} satisfy an “interlacing property” (that we will define exactly later on).
5. There exist interpolatory quadrature formulas

$$\int_{\mathbb{T}} F(z) d\mu(z) \approx \sum_{k=1}^n F(z_k^{(n)}) H_k^{(n)}, \quad (1)$$

where $(z_k^{(n)})_{k=1}^n$ are the zeros of F_n and $H_k^{(n)} > 0$, that are exact on certain $(2n - 1)$ -dimensional vector subspace of the space of Laurent polynomials, and that generate a sequence $(\mu_n)_{n \geq 1}$ of discrete measures supported on $\mathbb{T}$ and weakly converging to μ .

Notice that a SOP with respect to μ does not satisfy any of above properties. The search of these analog of quasi-orthogonal polynomials will lead to the so-called para-orthogonal polynomials ([9], [6], [3]). Finally, with a simple modification of these polynomials we will arrive at a sequence of Laurent polynomials called semi-orthogonal functions ([1], [2], [3]), that will provide the answer for the requirements 1, 2, 3, 4, 5.

Once we have found these sequences of functions, the final aim of this work will be to characterize the support of μ in terms of the asymptotic distribution of their zeros.

§ 2 - Basic tools.

We shall denote by $\mathbb{Z}$ the set of all integer numbers, and $\mathbb{N} = \{n \in \mathbb{Z} : n \geq 0\}$. Let $\mathbb{P}$ be the vector space of polynomials with complex coefficients, and $\mathbb{P}_n$ the vector subspace of polynomials whose degree is less than or equal to n .

Let μ be a probability measure supported on $\mathbb{T} := \{z \in \mathbb{C} : |z| = 1\}$. Let us define on $\mathbb{P}$ the inner product

$$\langle f, g \rangle := \int_{\mathbb{T}} f(z) \overline{g(z)} d\mu(z), \quad f, g \in \mathbb{P}.$$

The Gram-Schmidt procedure determines the sequence of monic orthogonal polynomials (SMOP), $(\Phi_n)_{n \in \mathbb{N}}$, satisfying the well known recurrence relation

$$\begin{aligned} \Phi_0(z) &= 1, \\ \Phi_{n+1}(z) &= z\Phi_n(z) + \Phi_{n+1}(0)\Phi_n^*(z), \quad n \in \mathbb{N}, \end{aligned}$$

where $\Phi_n^*(z) = z^n \overline{\Phi_n(z^{-1})}$ is the reversed polynomial of Φ_n . Furthermore, for $n \geq 1$, the zeros of the polynomial Φ_n lie on $|z| < 1$ and, thus, $|\Phi_n(0)| < 1$.

The sequence $(\Phi_n(0))_{n \in \mathbb{N}}$ is the so called sequence of Schur parameters, and the condition $|\Phi_n(0)| < 1$, $n \geq 1$ is in fact equivalent to the existence of an unique probability measure for which $(\Phi_n)_{n \in \mathbb{N}}$ is a SMOP (see [11], [7], [5]).

If we denote $e_n := \langle \Phi_n, \Phi_n \rangle > 0$, $n \geq 0$, then,

$$\begin{aligned} e_0 &= 1, \\ e_n &= \prod_{k=1}^n (1 - |\Phi_k(0)|^2), \quad n \geq 1, \end{aligned}$$

and $(e_n)_{n \geq 0}$ constitutes a strictly decreasing sequence. The orthonormal polynomial sequence related to above inner product, $(\varphi_n)_{n \geq 0}$, is $\varphi_n(z) = \frac{1}{\sqrt{e_n}} \Phi_n(z)$.

The Kernel polynomials $K_n(z, y)$ are defined by

$$K_n(z, y) := \sum_{k=0}^n \varphi_k(z) \overline{\varphi_k(y)} = \sum_{k=0}^n \frac{\Phi_k(z) \overline{\Phi_k(y)}}{e_k}, \quad n \in \mathbb{N}.$$

They satisfy the reproducing property

$$\langle K_n(z, y), f(z) \rangle = \overline{f(y)}, \quad f \in \mathbb{P}_n,$$

as well as the Christoffel-Darboux formula

$$e_{n+1}(1 - \overline{y}z)K_n(z, y) = \Phi_{n+1}^*(z) \overline{\Phi_{n+1}^*(y)} - \Phi_{n+1}(z) \overline{\Phi_{n+1}(y)}, \quad n \in \mathbb{N}.$$

Now, let us denote by $\Lambda = \text{span}\{z^n : n \in \mathbb{Z}\}$ the vector space of Laurent polynomials with complex coefficients, and, for $p, q \in \mathbb{Z}$ with $p \leq q$, let $\Lambda_{p,q}$ be the following vector subspace of Λ

$$\Lambda_{p,q} := \left\{ \sum_{k=p}^q \alpha_k z^k : \alpha_k \in \mathbb{C} \right\}.$$

The measure μ induces a linear functional on Λ , $\mathcal{L} : \Lambda \rightarrow \mathbb{C}$, given by

$$\mathcal{L}[f] := \int_{\mathbb{T}} f(z) d\mu(z), \quad f \in \Lambda.$$

An orthogonalization procedure on $(\Lambda_{-n,n}, \mathcal{L})$, $n \in \mathbb{N}$, leads to

$$\mathcal{B} = \{1\} \cup \left(\bigcup_{n \geq 1} \{f_n^{(1)}, f_n^{(2)}\} \right),$$

with

$$\left. \begin{aligned} f_n^{(1)}(z) &= \frac{\bar{\alpha}_n \Phi_{2n}(z) + \alpha_n \Phi_{2n}^*(z)}{z^n} = \frac{\bar{A}_n z \Phi_{2n-1}(z) + A_n \Phi_{2n-1}^*(z)}{z^n} \\ f_n^{(2)}(z) &= \frac{\bar{\beta}_n \Phi_{2n}(z) - \beta_n \Phi_{2n}^*(z)}{iz^n} = \frac{\bar{B}_n z \Phi_{2n-1}(z) - B_n \Phi_{2n-1}^*(z)}{iz^n} \end{aligned} \right\} \quad (2)$$

where $\alpha_n, \beta_n \in \mathbb{C}$ are such that $\operatorname{Re}(\alpha_n \bar{\beta}_n) \neq 0$. The set $\mathcal{B}$ of Laurent polynomials constitutes a basis for Λ such that

$$\mathcal{L}[f_n^{(j)}(z) z^k] = 0, \quad -n+1 \leq k \leq n-1, \quad j = 1, 2,$$

and the matrix

$$(\mathcal{L}[f_n^{(j)}(z) f_n^{(l)}(z)])_{j,l=1,2}$$

is positive definite for each $n \geq 1$. These functions $f_n^{(j)}$, $n \geq 1$, $j = 1, 2$, are called semi-orthogonal functions (SOF) with respect to $\mathcal{L}$ or μ (see [1], [2]).

Notice that, when $z \in \mathbb{T}$, then $f_n^{(j)}(z) \in \mathbb{R}$. So, SOF satisfy the first property we are looking for. To develop under which conditions they verify the rest of the requirements listed in the introduction, we start by studying the numerator polynomials of SOF.

Invariant para-orthogonal polynomials and quadrature formulas.

Definition. We say that $P \in \mathbb{P}_n$ is an n -para-orthogonal polynomial (n -POP) with respect to the measure μ if the following statements hold:

- i) $\langle P, z^k \rangle = 0, \quad 1 \leq k \leq n-1,$
- ii) $\langle P, 1 \rangle \neq 0, \quad \langle P, z^n \rangle \neq 0.$

Definition. For $\kappa \in \mathbb{C}$, $\kappa \neq 0$, a polynomial P is called (κ) -invariant if

$$P^*(z) = \kappa P(z), \quad \forall z \in \mathbb{C},$$

where $P^*(z) = z^{\deg(P)} \overline{P}(z^{-1})$.

Remark 1. Notice that $|P^*(z)| = |P(z)|$ for $z \in \mathbb{T}$ implies that necessarily $|\kappa| = 1$.

Proposition 1. A polynomial $P \in \mathbb{P}_n$ is an invariant n -POP if and only if there exist $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ such that $|\alpha| = |\beta|$ and

$$P(z) = \alpha \Phi_n(z) + \beta \Phi_n^*(z).$$

Proof. It follows immediately from the orthogonal decomposition

$$\mathbb{P}_n = z\mathbb{P}_{n-2} \oplus \text{span}\{\Phi_n, \Phi_n^*\}.$$

◇◇

Remark 2. Notice that the numerator of a SOF is a 1-invariant $2n$ -POP.

POP are introduced and studied for the first time in [9]. More recent results about them can be found in [6], [3]. Here, we will make only a summary of the mean properties that we need for our purposes (that is, properties relative to the zeros of POP), referring the reader to these works for the proofs.

Proposition 2. Invariant POP satisfy the following properties:

1. The zeros of an invariant n -POP are simple and lie on $\mathbb{T}$ (see [9], [3]).
2. Let P be a polynomial such that $P(z) = C_1 \Phi_n(z) + C_2 \Phi_n^*(z)$ and with zeros on $\mathbb{T}$. Then, P is an invariant n -POP (see [3]).
3. Let P be an invariant n -POP and let $z_0 \in \mathbb{T}$ be a zero of P . Then, we can write $P(z) = c(z - z_0)K_{n-1}(z, z_0)$, with $c \neq 0$. Conversely, $c(z - z_0)K_{n-1}(z, z_0)$ with $c \neq 0$ and $z_0 \in \mathbb{T}$ is an invariant n -POP (see [3]).
4. Let $P_n^{(1)}$ and $P_n^{(2)}$ be two invariant n -POP with respect to the same measure. If they have a common zero, then there exists $\lambda \in \mathbb{C} \setminus \{0\}$ such that $P_n^{(1)} = \lambda P_n^{(2)}$ (see [3]).
5. The vector subspace $\text{span}\{\Phi_n, \Phi_n^*\}$ is generated by any two invariant n -POP having not a common zero (see [3]).

Above properties are the starting point to construct interpolatory quadrature formulas like (1) that use as nodes the zeros of invariant POP. The results we summarize here can be found in [9]. Some extensions of them are discussed in [6], [3].

Given a measure μ supported on $\mathbb{T}$, we shall call an n -point Szegő quadrature problem, the following problem:

Find n points $(z_k)_{k=1}^n \subset \mathbb{T}$, $z_k \neq z_j$ if $k \neq j$ and n positive numbers $(H_k)_{k=1}^n$ such that the quadrature formula

$$\int_{\mathbb{T}} F(z) d\mu(z) \approx \sum_{k=1}^n F(z_k) H_k, \quad F \in C(\mathbb{T}), \quad (3)$$

(i) is exact for every $F \in \Lambda_{r,s}$, for some integers r, s ,

(ii) is not exact for every $F \in \Lambda_{r',s'}$ if $\Lambda_{r',s'} \not\supseteq \Lambda_{r,s}$.

Requirement (ii) is just a maximality condition, and is the responsible for the impossibility of ensuring the existence of a solution to above problem in general (see [3]). However, when $-r = s = n - 1$ such a maximal solution does always exists as the following theorem asserts.

Theorem 1. *The n -point Szegő quadrature problem has a solution on $\Lambda_{-n+1,n-1}$ if and only if the nodes $(z_k)_{k=1}^n$ are the zeros of an invariant n -POP (see [9], [6], [3]). In such a case, the weights H_k in the quadrature formula (3) are given by (see [6], [3])*

$$H_k = \frac{1}{K_{n-1}(z_k, z_k)}, \quad k = 1, \dots, n.$$

It is possible to find for the weights H_k another expression in terms of POP. Let $P(z)$ be an invariant n -POP whose zeros are $(z_k)_{k=1}^n$, and consider the rational function $f(z) = \frac{P(z)}{z^p}$ where $p \in \mathbb{Z}$ is fixed. Notice that $f \in \Lambda_{-p,n-p}$, and has the same zeros as P because $z = 0$ can not be a zero of P . Moreover, f provides a set of Laurent polynomials $(l_k)_{k=1}^n \subset \Lambda_{-p,n-p-1}$ through the following expressions

$$l_k(z) := \frac{f(z)}{(z - z_k)f'(z_k)}, \quad k = 1, \dots, n,$$

that is a basis for $\Lambda_{-p,n-p-1}$ because $l_k(z_j) = \delta_{kj}$. If $0 \leq p \leq n - 1$, then $\Lambda_{-p,n-p-1} \subset \Lambda_{-n+1,n-1}$, and the quadrature formula (3) that use as nodes the zeros of P is exact for l_k , $k = 1, \dots, n$. Therefore,

$$\int_{\mathbb{T}} l_k(z) d\mu(z) = \sum_{j=1}^n l_k(z_j) H_j = H_k, \quad k = 1, \dots, n.$$

Summarizing, if $P(z)$ is an invariant n -POP and p is an integer such that $0 \leq p \leq n - 1$, the rational function $f(z) = \frac{P(z)}{z^p}$ generate a solution to the n -point Szegő quadrature problem on $\Lambda_{-n+1,n-1}$: the nodes are the zeros of f and the weights are given in terms of f by

$$H_k = \int_{\mathbb{T}} \frac{f(z)}{(z - z_k)f'(z_k)} d\mu(z), \quad k = 1, \dots, n. \quad (4)$$

Of course, in spite of the p -dependence of f , the expression in the right hand side of (4) must be independent of p for $0 \leq p \leq n-1$ because the weight H_k so is.

If $\deg(P)$ is even, say $2n$, we can take $p = n$, and, thus

$$f(z) = \frac{\overline{\alpha}\Phi_{2n}(z) + \alpha\Phi_{2n}^*(z)}{z^n}.$$

In other words, $f(z)$ is a SOF with respect to μ and therefore real for $z \in \mathbb{T}$.

However, if $\deg(P) = 2n+1$, the choice for getting $f(z)$ real on $\mathbb{T}$ is $p = n + \frac{1}{2}$, and in this case

$$f(z) = \frac{\overline{\alpha}\Phi_{2n+1}(z) + \alpha\Phi_{2n+1}^*(z)}{z^n z^{\frac{1}{2}}}$$

is not a Laurent polynomial, and, even more, is not analytic in any open subset $\Omega \supset \mathbb{T}$ of $\mathbb{C}$. Moreover, the weights H_k can not be obtained like in (4) throughout such an f .

That is, if n is even, the quadrature formula can be generated by a SOF, but it is not such a case, in principle, when n is odd.

Study of the case odd n

Let

$$f(z) = \frac{\overline{\alpha}\Phi_{2n+1}(z) + \alpha\Phi_{2n+1}^*(z)}{z^n z^{\frac{1}{2}}},$$

where the determination for $z^{\frac{1}{2}}$ is arbitrarily chosen. We wish to find a SOF with the same zeros and that, therefore, can be used to generate quadrature formulas.

Obviously, the numerator of this f is a 1-invariant $(2n+1)$ -POP, and f vanishes at $2n+1$ different points on $\mathbb{T}$. Let w be one of them. Then, up to a non zero real factor,

$$f(z) = \frac{z-w}{iz^{\frac{1}{2}}w^{\frac{1}{2}}} \left(\frac{w}{z}\right)^n K_{2n}(z, w), \quad (5)$$

where again the determination of $w^{\frac{1}{2}}$ is arbitrarily chosen.

Let us consider the measure $\tilde{\mu}$, given by $d\tilde{\mu} = |z-w|^2 d\mu$ and denote $(\psi_n)_{n \in \mathbb{N}}$ the associated SMOP. Then, the sequences $(\Phi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$ are related by means of (see [8])

$$(z-w)\psi_{n-1}(z) = \Phi_n(z) - \frac{\Phi_n(w)}{K_{n-1}(w, w)} K_{n-1}(z, w).$$

Taking reversed polynomials, we get

$$-(z-w)\psi_{n-1}^*(z) = w\Phi_n^*(z) - \frac{\Phi_n^*(w)}{K_{n-1}(w, w)} z K_{n-1}(z, w).$$

Then

$$\begin{aligned}
(z-w)[\Phi_n^*(w)z\psi_{n-1}(z) + \Phi_n(w)\psi_{n-1}^*(z)] &= \\
&= \Phi_n^*(w)z\Phi_n(z) - w\Phi_n(w)\Phi_n^*(z) = \\
&= \frac{e_n}{e_{n-1}}[\Phi_{n+1}^*(w)\Phi_{n+1}(z) - \Phi_{n+1}(w)\Phi_{n+1}^*(z)]
\end{aligned}$$

follows from the elimination of $K_{n-1}(z, w)$ on above relationships and the recurrence formula. Thus, from (5),

$$f(z) = \frac{e_{2n+1}}{e_{2n}} \frac{z-w}{iz^{\frac{1}{2}}w^{\frac{1}{2}}} \frac{\Phi_{2n}^*(w)z\psi_{2n-1}(z) + \Phi_{2n}(w)\psi_{2n-1}^*(z)}{(zw)^n}.$$

Taking $z = e^{i\theta}$, $w = e^{i\beta}$, we have for $z \neq w$

$$\frac{z-w}{iz^{\frac{1}{2}}w^{\frac{1}{2}}} = \pm 2 \sin\left(\frac{\theta-\beta}{2}\right) \neq 0,$$

where the sign depends on the determination of $z^{\frac{1}{2}}$ and $w^{\frac{1}{2}}$. That is,

$$f(z) = \pm 2 \frac{e_{2n+1}}{e_{2n}} \sin\left(\frac{\theta-\beta}{2}\right) \frac{\Phi_{2n}^*(w)z\psi_{2n-1}(z) + \Phi_{2n}(w)\psi_{2n-1}^*(z)}{(zw)^n}. \quad (6)$$

So, if we define the Laurent polynomial

$$g(z) = \left(\frac{w}{z}\right)^n K_{2n}(z, w) \in \Lambda_{-n, n},$$

we can write, from (5) and (6)

$$g(z) = \frac{e_{2n+1}}{e_{2n}} \frac{\Phi_{2n}^*(w)z\psi_{2n-1}(z) + \Phi_{2n}(w)\psi_{2n-1}^*(z)}{(zw)^n}.$$

Notice that g has the same zeros as f , up to w . Moreover, as we can see from (2), g is a SOF with respect to $\tilde{\mu}$. Therefore, $g(z) \in \mathbb{R}$ for $z \in \mathbb{T}$, and its numerator is a 1-invariant $2n$ -POP with respect to $\tilde{\mu}$. Besides, $(z-w)g(z)$ is a Laurent polynomial which has the same zeros as f , and its numerator is a 1-invariant $(2n+1)$ -POP with respect to μ .

§ 3 - Semi-orthogonal functions and discrete measures.

For each $n \in \mathbb{N}$, we choose $w_n \in \mathbb{T}$. Then, we define a sequence $(F_n(z; w_n))_{n \geq 1}$ in Λ by means of

$$\left. \begin{aligned}
F_{2n}(z; w_{2n}) &:= \frac{\Phi_{2n}^*(w_{2n})\Phi_{2n}(z) - \Phi_{2n}(w_{2n})\Phi_{2n}^*(z)}{i(zw_{2n})^n}, \quad n \geq 1 \\
F_{2n+1}(z; w_{2n+1}) &:= \frac{\Phi_{2n}^*(w_{2n+1})z\psi_{2n-1}(z) + \Phi_{2n}(w_{2n+1})\psi_{2n-1}^*(z)}{(zw_{2n+1})^n}, \quad n \geq 0
\end{aligned} \right\} \quad (7)$$

so that $F_{2n}, F_{2n+1} \in \Lambda_{-n,n}$. This sequence is constituted by SOF: $(F_{2n})_{n \geq 1}$ is a sequence of SOF with respect to μ , and $(F_{2n+1})_{n \geq 0}$ is another one with respect to the family of varying measures $\mu^{(2n+1)}$, where

$$d\mu^{(2n+1)}(z) = |z - w_{2n+1}|^2 d\mu(z), \quad z \in \mathbb{T}.$$

Let us denote $(z_k^n)_{k=1}^n := \{z \in \mathbb{C} : F_n(z; w_n) = 0\} \cup \{w_n\}$. Now, the quadrature formulas generated by $F_{2n}(z; w_{2n})$ and $(z - w_{2n+1})F_{2n+1}(z; w_{2n+1})$ can be written together as

$$\int_{\mathbb{T}} F(z) d\mu(z) \approx \sum_{k=1}^n F(z_k^{(n)}) H_k^{(n)}, \quad (8)$$

where

$$H_k^{(n)} = \frac{1}{K_{n-1}(z_k^{(n)}, z_k^{(n)})}.$$

Moreover, formula (8) is exact for every $F \in \Lambda_{-n+1, n-1}$.

Above results can be read as follows. Let us consider the discrete measures on $\mathbb{T}$

$$d\mu_n(e^{i\theta}) = \sum_{k=1}^n \frac{1}{K_{n-1}(z_k^{(n)}, z_k^{(n)})} \delta(\theta - \theta_k^{(n)}) d\theta, \quad n \geq 1,$$

where $\theta_k^{(n)} = \text{Arg}(z_k^{(n)})$, and $(z_k^{(n)})_{k=1}^n$ are the zeros of

$$f_n(z; w_n) = \frac{\Phi_n^*(w_n)\Phi_n(z) - \Phi_n(w_n)\Phi_n^*(z)}{i(zw_n)^{\frac{n}{2}}}, \quad n \geq 1,$$

with $(w_n)_{n \geq 1} \subset \mathbb{T}$. Then, the exactness for $F \in \Lambda_{-n+1, n-1}$ means that

$$\int_{\mathbb{T}} F(z) d\mu_n(z) = \int_{\mathbb{T}} F(z) d\mu(z) \quad \forall F \in \Lambda_{-n+1, n-1}.$$

When $F = 1$ we get $\int_{\mathbb{T}} d\mu_n(z) = \int_{\mathbb{T}} d\mu(z) = c_0 = 1$, that is, $(\mu_n)_{n \geq 1}$ is a sequence of uniformly bounded measures. Therefore, for every $F \in C(\mathbb{T}), G \in \Lambda_{-m, m}$, if $n > m$ then

$$\begin{aligned} \left| \int_{\mathbb{T}} F(z) d\mu(z) - \int_{\mathbb{T}} F(z) d\mu_n(z) \right| &\leq \left| \int_{\mathbb{T}} (F(z) - G(z)) d\mu(z) \right| + \\ &\quad + \left| \int_{\mathbb{T}} (F(z) - G(z)) d\mu_n(z) \right| \leq 2\|F - G\|_{\infty}. \end{aligned}$$

Thus, since every continuous function on $\mathbb{T}$ can be uniformly approximated by Laurent polynomials,

$$\lim_n \int_{\mathbb{T}} F(z) d\mu_n(z) = \int_{\mathbb{T}} F(z) d\mu(z), \quad \forall F \in C(\mathbb{T}),$$

holds. That is, $\mu_n \rightarrow \mu$ in the $*$ -weak topology.

Summarizing, giving an arbitrary sequence $(w_n)_{n \geq 1} \subset \mathbb{T}$, we can define a sequence $(\mu_n)_{n \geq 1}$ of uniformly bounded discrete measures on $\mathbb{T}$ such that, μ_n is supported on the zeros of $f_n(z; w_n)$, and $\mu_n \xrightarrow{*} \mu$.

§ 4 - Sequences with interlacing zeros.

Let μ be a probability measure on $\mathbb{T}$ and let $(\Phi_n)_{n \in \mathbb{N}}$ be the related SMOP. For every sequence of complex numbers $(\alpha_n)_{n \geq 1}$, with $\alpha_n \neq 0$, we can define the functions

$$f_n(z) = \frac{\bar{\alpha}_n \Phi_n(z) - \alpha_n \Phi_n^*(z)}{iz^{\frac{n}{2}}}, \quad n \geq 1, \quad (9)$$

where the determination on $z^{\frac{n}{2}}$ can be arbitrarily chosen, because we are interested only in the location of their zeros. We know that f_n has their zeros on $\mathbb{T}$, $f_n(e^{i\theta}) \in \mathbb{R}$, and, moreover, a discrete measure μ_n is generated such that $\mu_n \xrightarrow{*} \mu$.

The study of the properties of the zeros of $(f_n)_{n \geq 1}$ is equivalent to the study of the same properties for the SOF $(F_n(z; w_n))_{n \geq 1}$ given in (7).

By using the recurrence relation for $(\Phi_n)_{n \in \mathbb{N}}$ in (9), we have

$$f_n(z) = \frac{e_n}{e_{n-1}} \frac{\bar{\beta}_n z \Phi_{n-1}(z) - \beta_n \Phi_{n-1}^*(z)}{iz^{\frac{n}{2}}}, \quad n \geq 1, \quad (10)$$

where $\beta_n = \frac{e_{n-1}}{e_n}(\alpha_n - \Phi_n(0)\bar{\alpha}_n)$, $\bar{\beta}_n = \frac{e_{n-1}}{e_n}(\bar{\alpha}_n - \overline{\Phi_n(0)\alpha_n})$, and thus

$$\alpha_n = \beta_n + \Phi_n(0)\bar{\beta}_n, \quad n \geq 1. \quad (11)$$

Without loss of generality, we can assume that $(\alpha_n)_{n \geq 1}$ is given by means of a polynomial sequence $(p_n)_{n \geq 1}$ in the following way

$$\alpha_n := w^{-\frac{m}{2}} p_n(w),$$

where $w \in \mathbb{T}$ is arbitrary, and $m = m(n) = \deg(p_n)$, which is chosen so that $m(n+1) = m(n) + 1$. Under these assumptions, we can write

$$\beta_n = w^{-\frac{m}{2}} \frac{e_{n-1}}{e_n} [p_n(w) - \Phi_n(0)p_n^*(w)].$$

Let $\omega_0 \in \mathbb{R}$ be fixed. We can write $\mathbb{T} = \{e^{i\theta} : \theta \in [\omega_0, \omega_0 + 2\pi)\}$, and, thus, $f_n(e^{i\theta})$ is a C^∞ real function defined on $[\omega_0, \omega_0 + 2\pi)$, where the non integer powers

of z and w are taken so that $\arg(z^{\frac{1}{2}}), \arg(w^{\frac{1}{2}}) \in [\frac{\omega_0}{2}, \frac{\omega_0}{2} + \pi)$. For two numbers $\zeta_1, \zeta_2 \in \mathbb{T}$, let us denote $\theta_j = \arg(\zeta_j) \in [\omega_0, \omega_0 + 2\pi)$. Then, we can establish an order relation by

$$\zeta_1 < \zeta_2 \iff \theta_1 < \theta_2.$$

Let $(\zeta_j^{(n)})_{j=1}^n$ and $(\zeta_j^{(n+1)})_{j=1}^{n+1}$ be the zeros of f_n and f_{n+1} respectively, which are ordered as above. Then, we wish to determine a suitable sequence $(\alpha_n)_{n \geq 1}$ such that, for each $n \geq 1$,

$$\zeta_j^{(n+1)} < \zeta_j^{(n)} < \zeta_{j+1}^{(n+1)}, \quad j = 1, \dots, n,$$

holds when $\zeta_j^{(k)} \neq e^{i\omega_0}$, $k = n, n+1$. In short, in such a case we say that f_n and f_{n+1} have interlacing zeros. It is a well known result, derived from Sturm's theorem, that the interlacing condition is equivalent to

$$\operatorname{sgn}\left[\left(\frac{d}{d\theta} f_{n+1}(\zeta)\right) f_n(\zeta)\right] = \operatorname{sgn}\left[i\zeta f'_{n+1}(\zeta) f_n(\zeta)\right] = \text{constant}$$

for all $\zeta \in (\zeta_j^{(n+1)})_{j=1}^{n+1}$, $\zeta \neq e^{i\omega_0}$.

Let us consider for f_n and f_{n+1} the expressions (9) and (10), respectively,

$$\begin{aligned} f_n(z) &= \frac{\bar{\alpha}_n \Phi_n(z) - \alpha_n \Phi_n^*(z)}{iz^{\frac{n}{2}}}, \\ f_{n+1}(z) &= \frac{e_{n+1}}{e_n} \frac{\bar{\beta}_{n+1} z \Phi_n(z) - \beta_{n+1} \Phi_n^*(z)}{iz^{\frac{n+1}{2}}}. \end{aligned}$$

Since $f_{n+1}(\zeta) = 0$, we have $\frac{\beta_{n+1}}{\bar{\beta}_{n+1}} = \frac{\zeta \Phi_n(\zeta)}{\Phi_n^*(\zeta)}$. Thus

$$\beta_{n+1} = k_{n+1} \zeta^{\frac{-n+1}{2}} \Phi_n(\zeta), \quad \bar{\beta}_{n+1} = k_{n+1} \zeta^{\frac{-n-1}{2}} \Phi_n^*(\zeta), \quad (12)$$

with $k_{n+1} = k_{n+1}(\zeta) \in \mathbb{R} \setminus \{0\}$. For f_{n+1} , and using Christoffel-Darboux formula, we get

$$f_{n+1}(z) = k_{n+1} e_{n+1} \frac{z - \zeta}{i\zeta} \left(\frac{\zeta}{z}\right)^{\frac{n}{2}} K_n(z, \zeta).$$

Hence,

$$i\zeta f'_{n+1}(\zeta) = k_{n+1} e_{n+1} K_n(\zeta, \zeta).$$

On the other hand, from (12),

$$f_n(\zeta) = \frac{1}{k_{n+1}} \frac{\bar{\alpha}_n \beta_{n+1} \zeta^{-\frac{1}{2}} - \alpha_n \bar{\beta}_{n+1} \zeta^{\frac{1}{2}}}{i}$$

follows. That is,

$$i\zeta f'_{n+1}(\zeta)f_n(\zeta) = 2e_{n+1}K_n(\zeta, \zeta) \frac{\overline{\alpha}_n\beta_{n+1}\zeta^{-\frac{1}{2}} - \alpha_n\overline{\beta}_{n+1}\zeta^{\frac{1}{2}}}{2i}.$$

The last factor in the right hand side suggests the choice

$$w = e^{i\omega_0}, \quad \beta_{n+1} = w^{\frac{1}{2}}\alpha_n, \quad (13)$$

that leads to an expression with constant sign

$$\frac{\overline{\alpha}_n\beta_{n+1}\zeta^{-\frac{1}{2}} - \alpha_n\overline{\beta}_{n+1}\zeta^{\frac{1}{2}}}{2i} = |\alpha_n|^2 \sin\left(\frac{\omega_0 - \theta}{2}\right) < 0,$$

where $\theta = \arg(\zeta) \in (\omega_0, \omega_0 + 2\pi)$.

Now, from (13) and (11) it follows that

$$p_{n+1}(w) = wp_n(w) + \Phi_{n+1}(0)p_n^*(w), \quad n \geq 1. \quad (14)$$

This equation has two independent solutions (see [7], [5]), that can be chosen as the sequences $(\Phi_n(w))_{n \geq 1}$ and $(i\Omega_n(w))_{n \geq 1}$, where Ω_n is the n -th second kind polynomial. Thus, there exist two fixed polynomials A , B , and an integer $k \geq 0$, such that

$$\begin{cases} p_n(w) = A(w)\Phi_n(w) + B(w)\Omega_n(w), \\ A^{*k}(w) = A(w), \quad B^{*k}(w) = -B(w), \end{cases} \quad (15)$$

where $P^{*k}(z) = z^k \overline{P}(z^{-1})$, $\deg(P) \leq k$. The conditions (14) and (15) are equivalents (see [10]).

Summarizing, we can deduce the following result.

Theorem 2. *Let $(p_n)_{n \in \mathbb{N}}$ be a polynomial sequence satisfying (14), or equivalently (15). Let $w = e^{i\omega_0}$ be fixed. Then, for the sequence of functions $(f_n)_{n \geq 1}$ defined by*

$$f_n(z) = \frac{\overline{p_n(w)}\Phi_n(z) - p_n(w)\Phi_n^*(z)}{iz^{\frac{n}{2}}}, \quad n \geq 1, \quad (16)$$

f_n and f_{n+1} have interlacing zeros in $(\omega_0, \omega_0 + 2\pi)$.

As two particular cases of (16), we have

$$\begin{aligned} f_n^{(1)}(z; w) &:= \frac{\Phi_n^*(w)\Phi_n(z) - \Phi_n(w)\Phi_n^*(z)}{i(zw)^{\frac{n}{2}}}, & n \geq 1, \\ f_n^{(2)}(z; w) &:= \frac{\Omega_n^*(w)\Phi_n(z) + \Omega_n(w)\Phi_n^*(z)}{(zw)^{\frac{n}{2}}}, & n \geq 1. \end{aligned} \quad (17)$$

From (15) and (16),

$$\begin{aligned} f_n(z) &= \frac{A(w)}{w^{\frac{k}{2}}} f_n^{(1)}(z; w) - \frac{B(w)}{w^{\frac{k}{2}}} f_n^{(2)}(z; w) = \\ &= A_1 f_n^{(1)}(z; w) + A_2 f_n^{(2)}(z; w), \end{aligned}$$

(where $A_1 = \frac{A(w)}{w^{\frac{k}{2}}}$, $A_2 = -\frac{B(w)}{w^{\frac{k}{2}}}$) follows. Moreover, A_1 and A_2 are real numbers, independent from n , with $(A_1, A_2) \neq (0, 0)$.

Remark 3. Conversely, given $f_n(z) = \frac{\overline{\alpha}_n \Phi_n(z) - \alpha_n \Phi_n^*(z)}{iz^{\frac{n}{2}}}$, $\alpha_n \neq 0$, there exist two real numbers $A_1(n)$, $A_2(n)$ with $(A_1(n), A_2(n)) \neq (0, 0)$ such that

$$f_n(z) = A_1(n) f_n^{(1)}(z; w) + A_2(n) f_n^{(2)}(z; w).$$

In fact, let us denote

$$\begin{aligned} P_n(z) &= \overline{\alpha}_n \Phi_n(z) - \alpha_n \Phi_n^*(z), \\ P_n^{(1)}(z) &= w^{-\frac{n}{2}} [\Phi_n^*(w) \Phi_n(z) - \Phi_n(w) \Phi_n^*(z)], \\ P_n^{(2)}(z) &= w^{-\frac{n}{2}} [\Omega_n^*(w) \Phi_n(z) + \Omega_n(w) \Phi_n^*(z)]. \end{aligned}$$

These polynomials are 1-invariant n -POP. Moreover,

$$P_n \in \text{span}\{\Phi_n, \Phi_n^*\} = \text{span}\{P_n^{(1)}, P_n^{(2)}\}.$$

Thus, there exists $(A_1(n), A_2(n)) \neq (0, 0)$ such that

$$P_n(z) = A_1(n) P_n^{(1)}(z) + \frac{A_2(n)}{i} P_n^{(2)}(z)$$

holds. Thus,

$$f_n(z) = A_1(n) f_n^{(1)}(z; w) + A_2(n) f_n^{(2)}(z; w).$$

By taking $z = w$ and $z \in \mathbb{T}$, it follows that $A_1(n), A_2(n) \in \mathbb{R}$.

The choice (13), which guarantees interlacing property, corresponds to the case when $A_1(n)$, $A_2(n)$ are independent of n . Of course, (13) is not the unique option for interlacing zeros.

Remark 4. Notice that the functions $f_n^{(1)}(z; w)$ have a common zero at $\zeta_1^{(n)} = w$. In this case, the interlacing property is strictly verified in the interval $(\omega_0, \omega_0 + 2\pi)$. By using the well-known identity

$$\Omega_n^*(z) \Phi_n(z) + \Omega_n(z) \Phi_n^*(z) = 2e_n z^n \tag{18}$$

(see [7]), we obtain that $f_n^{(2)}(w; w) = 2e_n$. This implies that, for the remaining cases, i.e., when $B(w) \neq 0$ in (15), w is not a zero of f_n . Thus, the interlacing property holds in $[\omega_0, \omega_0 + 2\pi)$.

Finally, we are able to give other kind of interlacing properties.

Proposition 3. *For each $n \geq 1$, the functions $f_n^{(1)}(z; w)$ and $f_n^{(2)}(z; w)$ have interlacing zeros in $[\omega_0, \omega_0 + 2\pi)$.*

Proof. It is enough to prove that the expression

$$i\zeta f_n^{(1)'}(\zeta; w) f_n^{(2)}(\zeta; w)$$

has the same sign for every zero ζ of $f_n^{(1)}(z; w)$. If $f_n^{(1)}(\zeta; w) = 0$, then

$$\frac{\Phi_n(\zeta)}{\Phi_n(w)} = \frac{\Phi_n^*(\zeta)}{\Phi_n^*(w)}.$$

Using (18) we get

$$\begin{aligned} f_n^{(2)}(\zeta; w) &= \frac{\Omega_n^*(w)\Phi_n(w) + \Omega_n(w)\Phi_n^*(w)}{(\zeta w)^{\frac{n}{2}}} \frac{\Phi_n^*(\zeta)}{\Phi_n^*(w)} = \\ &= 2e_n \frac{\Phi_n^*(\zeta)}{\Phi_n^*(w)} \left(\frac{w}{\zeta} \right)^{\frac{n}{2}}. \end{aligned}$$

On the other hand,

$$\begin{aligned} f_n^{(1)}(z; w) &= \frac{\Phi_n^*(w)}{\Phi_n^*(\zeta)} \left(\frac{\zeta}{w} \right)^{\frac{n}{2}} \frac{\Phi_n^*(\zeta)\Phi_n(z) - \Phi_n(\zeta)\Phi_n^*(z)}{i(z\zeta)^{\frac{n}{2}}} = \\ &= \frac{\Phi_n^*(w)}{\Phi_n^*(\zeta)} \left(\frac{\zeta}{w} \right)^{\frac{n}{2}} f_n^{(1)}(z; \zeta) = \\ &= \frac{\Phi_n^*(w)}{\Phi_n^*(\zeta)} \frac{\zeta}{(wz)^{\frac{n}{2}}} e_n K_{n-1}(z, \zeta) \frac{z - \zeta}{i\zeta}. \end{aligned}$$

Thus,

$$i\zeta f_n^{(1)'}(\zeta; w) = \frac{\Phi_n^*(w)}{\Phi_n^*(\zeta)} \left(\frac{\zeta}{w} \right)^{\frac{n}{2}} e_n K_{n-1}(\zeta, \zeta).$$

Finally,

$$i\zeta f_n^{(1)'}(\zeta; w) f_n^{(2)}(\zeta; w) = 2e_n^2 K_{n-1}(\zeta, \zeta) > 0$$

for each zero ζ of $f_n^{(1)}(z; w)$. $\diamond\diamond$

Corollary 1. *Let $A_j, B_j \in \mathbb{R}$, $j = 1, 2$, and*

$$\begin{aligned} f_n(z) &= A_1 f_n^{(1)}(z; w) + A_2 f_n^{(2)}(z; w), \\ g_n(z) &= B_1 f_n^{(1)}(z; w) + B_2 f_n^{(2)}(z; w), \end{aligned}$$

where $\begin{vmatrix} A_1 & A_2 \\ B_1 & B_2 \end{vmatrix} \neq 0$. Then, f_n and g_n have interlacing zeros in $[\omega_0, \omega_0 + 2\pi)$.

Proof. Notice that f_n and g_n are linearly independent. Then, from Proposition 2, they have not a common zero. Let ζ_0 be such that $f_n(\zeta_0) = 0$. Then, up to a non-zero real factor, $f_n(z) = f_n^{(1)}(z; \zeta_0)$ holds. Thus, again from Proposition 2, we have that there exist $C_1, C_2 \in \mathbb{R}$, with $C_2 \neq 0$, such that $g_n(z) = C_1 f_n^{(1)}(z; \zeta_0) + C_2 f_n^{(2)}(z; \zeta_0)$. Hence, by using Proposition 3, we have for each zero ζ of f_n that

$$\begin{aligned} i\zeta f_n'(\zeta)g_n(\zeta) &= i\zeta f_n^{(1)'}(\zeta; \zeta_0)C_2 f_n^{(2)}(\zeta; \zeta_0) = \\ &= 2e_n^2 K_{n-1}(\zeta, \zeta)C_2, \end{aligned}$$

which keeps constant sign. $\diamond\diamond$

§ 6 - Semi-orthogonal functions and support of the orthogonality measure.

We have find that any sequence of functions given in Theorem 2 satisfies properties 1, 2, 4, 5 listed in §1. It remains to reach the property 3, as well as to develop the relation between $\text{supp } \mu$ and the asymptotic distribution of zeros of such kind of functions. The starting point for this relation are the quadrature formulas described in §2, that are constructed by using as nodes the zeros of functions with the general form

$$f_n(z) = \frac{\overline{\alpha}_n \Phi_n(z) - \alpha_n \Phi_n^*(z)}{iz^{\frac{n}{2}}}, \quad \alpha_n \neq 0. \quad (19)$$

As we said in §3, these quadrature formulas provide discrete measures μ_n weakly converging to the orthogonality measure μ , and supported on the zeros of the functions f_n . As a consequence, given an arbitrary sequence $(f_n)_{n \geq 1}$ of functions with the form (19), a point of $\text{supp } \mu$ must be a limit point of the set of zeros of all the functions f_n , or a zero of infinitely many functions f_n . Thus, denoting by A' the derived set of A , we have the following proposition.

Proposition 4. *Let $(f_n)_{n \geq 1}$ be any sequence of functions with the form (19), and let $\mathcal{Z}$ and $\mathcal{X}$ be the set of zeros of all the functions f_n , and the set of complex numbers that are zeros of infinitely many functions f_n , respectively. Then,*

$$\text{supp } \mu \subseteq \mathcal{Z}' \cup \mathcal{X}.$$

Since $\mathcal{Z}'' \subset \mathcal{Z}'$ and $\mathcal{X}' \subset \mathcal{Z}'$, a corollary follows immediately.

Corollary 2. *Under the assumptions of Proposition 4,*

$$(\text{supp } \mu)' \subseteq \mathcal{Z}'.$$

Corollary 2 says that a limit point of $\text{supp } \mu$ must be a limit point of $\mathcal{Z}$ too. From Proposition 4, the rest of the points of $\text{supp } \mu$, that is, the isolated points, could be limit points of $\mathcal{Z}$, or zeros of infinitely many functions f_n . Now, we are going to demonstrate that, indeed, an isolated point z_0 of $\text{supp } \mu$ that is not a limit point of $\mathcal{Z}$, must be a zero of all the functions f_n except, at most, finitely many of them. If it were not the case, then there would be a subsequence $(f_{n_k})_{k \in \mathbb{N}}$ such that $f_{n_k}(z_0) \neq 0$ for $k \in \mathbb{N}$. Moreover, we are supposing that z_0 is an isolated point of $\text{supp } \mu$ that is not a limit point of $\mathcal{Z}$, so, there must be a neighborhood $\mathcal{U}$ of z_0 where none of the functions f_{n_k} can vanish, and such that $\mathcal{U} \cap \text{supp } \mu = \{z_0\}$. Therefore, $\int_{\mathcal{U} \cap \mathbb{T}} d\mu = \mu(\{z_0\}) \neq 0$. Now, let us consider the subsequence of measures $(\mu_{n_k})_{k \in \mathbb{N}}$ corresponding to the subsequence of functions $(f_{n_k})_{k \in \mathbb{N}}$. Since $(\mu_n)_{n \in \mathbb{N}}$ is uniformly bounded, $(\mu_{n_k})_{k \in \mathbb{N}}$ so is, and Helly's theorem implies the existence of a subsubsequence $(\mu_{n_{k_j}})_{j \in \mathbb{N}}$ which is convergent to μ a.e.. For this subsubsequence it must be

$$\lim_j \int_{\mathcal{U} \cap \mathbb{T}} d\mu_{n_{k_j}} = \int_{\mathcal{U} \cap \mathbb{T}} d\mu.$$

But, since the functions f_{n_k} have no zeros in $\mathcal{U}$, it is $\mathcal{U} \cap \text{supp } \mu_{n_{k_j}} = \emptyset$, and, hence, $\int_{\mathcal{U} \cap \mathbb{T}} d\mu_{n_{k_j}} = 0$, which is in contradiction with $\int_{\mathcal{U} \cap \mathbb{T}} d\mu \neq 0$.

Then, we can state the following result.

Proposition 5. *Under the assumptions of Proposition 4,*

$$\text{supp } \mu \subseteq \mathcal{Z}' \cup \tilde{\mathcal{X}},$$

where $\tilde{\mathcal{X}}$ is the set of complex numbers that are zeros of all the functions f_n except, at most, finitely many of them.

To go beyond these results we need a property that is quasi-reciprocal of previous one, and, therefore, leads to an inclusion in opposite sense. Proposition 5 says that if a connected subset of $\mathbb{T}$ is “eventually free of zeros”, then its interior is “out of the support of μ ”. On the contrary, the following proposition implies that if a connected subset of $\mathbb{T}$ is “out of the support of μ ”, then its closure contains at most one zero.

Proposition 6. *Let $(f_n)_{n \geq 1}$ be any sequence of functions with the form (19), and let z_1, z_2 be two zeros of f_n . Then, in any of the two connected components of $\mathbb{T} \setminus \{z_1, z_2\}$ there exists at least one point of $\text{supp } \mu$, as well as one zero of f_m for $m > n$.*

Proof. Since we are just interested in the location of zeros we will only work with the polynomials $P_n(z) = \bar{\alpha}_n \Phi_n(z) - \alpha_n \Phi_n^*(z)$ that have the same zeros as f_n . Notice first that $\frac{zP_n(z)}{(z-z_1)(z-z_2)} \in \text{span}\{1, z, z^2, \dots, z^{n-1}\}$, and, therefore,

$$\int_{\mathbb{T}} \frac{zP_n(z)}{(z-z_1)(z-z_2)} \bar{P}_n(z^{-1}) d\mu(z) = 0,$$

due to the orthogonality properties of Φ_n and Φ_n^* . If $\theta_j = \text{Arg}(z_j)$, above result reads

$$\int_0^{2\pi} \frac{|P_n(e^{i\theta})|^2}{\sin(\frac{\theta-\theta_1}{2}) \sin(\frac{\theta-\theta_2}{2})} d\mu(e^{i\theta}) = 0.$$

But the expression inside the integral has now constant and opposite sign in each connected component of $\mathbb{T} \setminus \{z_1, z_2\}$, except for the zeros of f_n , for which is null (including z_1 and z_2). Thus, taking into account that $\text{supp } \mu$ is an infinite set, if the integral has to vanish, then there must be points of $\text{supp } \mu$ in both components.

On the other hand, we have that $\frac{zP_n(z)}{(z-z_1)(z-z_2)} \bar{P}_n(z^{-1}) \in \Lambda_{-n+1, n-1}$, so, the quadrature formulas described in §3 are exact over this Laurent polynomial if we choose as nodes the zeros $z_k^{(m)}$ of f_m for $m > n$. Hence, if $\theta_k^{(m)} = \text{Arg}(z_k^{(m)})$, it must be

$$\sum_{k=1}^m \frac{|P_n(e^{i\theta_k^{(m)}})|^2}{\sin(\frac{\theta_k^{(m)}-\theta_1}{2}) \sin(\frac{\theta_k^{(m)}-\theta_2}{2})} H_k^{(m)} = 0, \quad m > n.$$

If all the zeros $z_k^{(m)}$ were located in only one of the connected components of $\mathbb{T} \setminus \{z_1, z_2\}$, all the non-null terms in this sum would have the same sign. Since there must be non-null terms because the number of zeros of P_n is $n < m$, this is in contradiction with the fact that above sum must vanish. $\diamond$

Notice that this proposition has as a direct consequence the property 3 listed in §1 that we were looking for. That is,

Corollary 3. *If I is a connected subset of $\mathbb{T}$ such that $\text{supp } \mu \subseteq I$, at most one zero of f_n lies on the closure of $\mathbb{T} \setminus I$.*

To get more accurate results about $\text{supp } \mu$ we need to restrict our attention to those sequences of functions with interlacing zeros. Then, we will suppose that $(f_n(z))_{n \geq 1}$ is one of the sequences described in Theorem 2, that is,

$$f_n(z) = A_1 f_n^{(1)}(z; w) + A_2 f_n^{(2)}(z; w), \quad (A_1, A_2) \neq (0, 0), \quad (20)$$

where $f_n^{(i)}$, $i = 1, 2$, are given in (17). In that case, f_n and f_{n+1} can have only one common zero at the point w , due to the interlacing of zeros in $\mathbb{T} \setminus \{w\}$. Thus,

$\tilde{\mathcal{X}} \subseteq \{w\}$. In fact, from Remark 4, we can see that $\tilde{\mathcal{X}} = \{w\}$ only when $A_2 = 0$. Thus, we have reached the following result.

Proposition 7. *Let $(f_n)_{n \geq 1}$ be any sequence of functions with the form (20). Then, if $A_2 = 0$,*

$$\text{supp } \mu \subseteq \mathcal{Z}' \cup \{w\},$$

and, if $A_2 \neq 0$,

$$\text{supp } \mu \subseteq \mathcal{Z}'.$$

Now, we will work only with the functions $f_n^{(1)}(z; w)$, that is, those functions $f_n(z)$ with a common zero at $z = w$. We will suppose for $\text{supp } \mu$ a non-trivial situation, what means that $\text{supp } \mu \neq \mathbb{T}$. The strategy will be to choose a point w in each connected component of $\mathbb{T} \setminus \text{supp } \mu$, and consider the corresponding functions $f_n^{(1)}(z; w)$. So that, the representation of $\text{supp } \mu$ will be given throughout the zeros of several sequences $(f_n^{(1)}(z; w))_{n \geq 1}$, one for each connected component of $\mathbb{T} \setminus \text{supp } \mu$.

Theorem 3. *Suppose that $\text{supp } \mu \neq \mathbb{T}$ and let*

$$\mathbb{T} \setminus \text{supp } \mu = \bigcup_{k \in \mathcal{A}} C_k$$

be the decomposition of $\mathbb{T} \setminus \text{supp } \mu$ into connected components. If $w_k \in C_k$ for every $k \in \mathcal{A}$, then,

$$\text{supp } \mu = \bigcap_{k \in \mathcal{A}} \mathcal{Z}(w_k)',$$

where $\mathcal{Z}(w)$ is the set of zeros of all the functions $f_n^{(1)}(z; w)$.

Proof. From Proposition 7 it is clear that, since $w_k \notin \text{supp } \mu$, it has to be $\text{supp } \mu \subseteq \bigcap_{k \in \mathcal{A}} \mathcal{Z}(w_k)'$. On the other hand, Proposition 6 implies that w_k must be the unique zero of $f_n^{(1)}(z; w_k)$ in C_k . Therefore, $\mathcal{Z}(w_k)' \cap C_k = \emptyset$, and, thus, $\bigcap_{k \in \mathcal{A}} \mathcal{Z}(w_k)' \subseteq \text{supp } \mu$. $\diamond\diamond$

Acknowledgements.- This research was partially supported by Universidad de Zaragoza PAI UZ-97-CIE-10 and Dirección General de Enseñanza superior of Spain (DGES) Project PB98-1615.

References

- [1] ALFARO, M.; CANTERO, M.J.; MORAL, L. "Semi-orthogonal functions and orthogonal polynomials on the unit circle", J. Comput. Appl. Math. **99** (1998) 3-14.
- [2] CANTERO, M.J. "Polinomios ortogonales sobre la circunferencia unidad. Modificaciones de los parámetros de Schur", Doctoral Dissertation, Universidad de Zaragoza, 1997.
- [3] CANTERO, M.J.; FERRER, M.P.; MORAL, L. ; VELAZQUEZ, L. "Funciones semiortogonales y formulas de cuadratura", Actes des *VI^{èmes}* Journées de Mathématiques Appliquées et de Statistiques (Jaca, 1999) pp. 141–150, Publications de l'Université de Pau, 2001.
- [4] CHIHARA, T.S. "An introduction to orthogonal polynomials", Gordon and Breach, New York, 1978.
- [5] FREUD, G. "Orthogonal polynomials", Pergamon Press, Oxford, 1971.
- [6] BULTHEEL, A; GONZÁLEZ-VERA, P; HENDRIKSEN, E; NJÄSTAD, O. "Orthogonal rational functions", University Press, Cambridge, 1999.
- [7] GERONIMUS, Ya.L. "Orthogonal polynomials", Consultants Bureau, New York, 1961.
- [8] GODOY, E; MARCELLAN, F. "Orthogonal polynomials and rational modifications of measures", Canad. J. Math. **45** (1993) 930-943.
- [9] JONES, W.B.; NJÄSTAD, O; THRON, W.I. "Moment theory, orthogonal polynomials, quadrature, and continued fractions associated with the unit circle", Bull. London Math. Soc. **21** (1989) 113-152.
- [10] PEHERSTORFER, F; STEINBAUER, R. "Characterization of general orthogonal polynomials with respect to a functional", J. Comp. Appl. Math. **65** (1995) 339-355.
- [11] SZEGŐ, G. "Orthogonal Polynomials", Amer. Math. Soc. Colloq. Publ. 23, Providence, Rhode Island, 1975 (Fourth Edition).